Author Index

Appleton, J. D., see Cook, A. P. 395
Aue, W. A., see Sun, X.-Y. 75
Azoulay, M., see Morineau, G. 209
Baczynskyj, W. M., see Richfield-Fratz, N. 167
Baer, J. C.
——, Slack, J. A. and Pettit, G. R.
Stability-indicating high-performance liquid chromatography assay for the anticancer drug bryostatin 1 332
Baeyens, W., see Xianren, Q. 15
Bailey, J. E., Jr., see Richfield-Fratz, N. 167
Barry, E. F., see Costanzo, R. B. 373
Bartle, K. D., see Raynor, M. W. 292
Belenkii, A. B., see Ruban, V. F. 41
Belenkii, B. G., see Ruban, V. F. 41
Betts, T. J.
Use of an inexpensive low-temperature liquid crystal stationary phase for the study of volatile oil constituents with short gas chromatographic retention times 272
Bianchi, A.
——, Varney, M. S. and Phillips, J.
Modified analytical technique for the determination of trace organics in water using dynamic headspace and gas chromatography–mass spectrometry 111
Biondi, P. A.
——, Manca, F., Negri, A., Tedeschi, G. and Secchi, C.
Gas chromatographic determination of glycoprotein amino sugars as O-pentafluorobenzyloxime acetates 315
Bitetto, M. di, see Gobetti, M. 327
Blakemore, W. M., see Thompson, H. C., Jr. 159
Blink, A.
——, Suijkerbuijk, M. L., Ishiwata, T. and Feringa, B. L.
Gas chromatographic method for enantiomeric excess determination of alcohols not requiring chiral auxiliary compounds or chiral stationary phases 285
Block, G. G. A. G., see Delcour, J. A. 149
Böhland, H., see Radicke, J. 304
Boreham, C. J.
—— and Fookes, C. J. R.
Separation of nickel(II) alkylporphyrins by reversed-phase high-performance liquid chromatography. Methodology and application 195
Boter, H. L., see Kientz, Ch. E. 385
Božnar, B., see Žmitek, J. 276
Brinkman, U. A. Th., see Kientz, Ch. E. 385
Büyüktimkin, N.
——, Keller, F. and Schunack, W.
Gas–liquid chromatographic resolution of some racemic synthons for lamtidine analogous histamine H_2-receptor antagonists via diastereomeric amides of (1*S*)-(−)-camphanic acid 402
Caude, M., see Lienne, M. 406
——, see Macaudière, P. 357
——, see Nizery, D. 49
Chan, P. F., see Law, B. 267
Cheah, I. C. L.
——, Sitaram, B. R., Pappas, A., Thi, N. L., Finnin, B. C. and Reed, B. L.
Normal-phase and reversed-phase liquid chromatographic techniques for the determination of dithranol and its degradation products 414
Chornet, E., see Thring, R. W. 441
Cook, A. P.
——, Macleod, T. M., Appleton, J. D. and Fell, A. F.
Reversed-phase high-performance liquid chromatographic method for the quantification of 5-hydroxymethylfurfural as the major degradation product of glucose in infusion fluids 395
Costanzo, R. B.
—— and Barry, E. F.
Gas chromatographic detection of selected organochlorine species using an alternating current plasma detector 373
De Jong, G. J., see Kientz, Ch. E. 385
Dekker, M., see Griest, W. H. 423
Delcour, J. A.
——, Vinkx, C. J. A., Vanhamel, S. and Block, G. G. A. G.
Combined monitoring of UV absorbance and fluorescence intensity as a diagnostic criterion in reversed-phase high-performance liquid chromatographic separation of natural phenolic acids 149
De Pozuelo, M. M., see Reuvers, Th. B. A. 321
Di Bitetto, M., see Gobetti, M. 327
Dreux, M., see Nizery, D. 49
Ellegren, H.
—— and Låås, T.
Size-exclusion chromatography of DNA restriction fragments. Fragment length determinations and a comparison with the behaviour of proteins in size-exclusion chromatography 217

Author Index

Appleton, J. D., see Cook, A. P. 395
Aue, W. A., see Sun, X.-Y. 75
Azoulay, M., see Morineau, G. 209
Baczynskyj, W. M., see Richfield-Fratz, N. 167
Baer, J. C.
——, Slack, J. A. and Pettit, G. R.
Stability-indicating high-performance liquid chromatography assay for the anticancer drug bryostatin 1 332
Baeyens, W., see Xianren, Q. 15
Bailey, J. E., Jr., see Richfield-Fratz, N. 167
Barry, E. F., see Costanzo, R. B. 373
Bartle, K. D., see Raynor, M. W. 292
Belenkii, A. B., see Ruban, V. F. 41
Belenkii, B. G., see Ruban, V. F. 41
Betts, T. J.
Use of an inexpensive low-temperature liquid crystal stationary phase for the study of volatile oil constituents with short gas chromatographic retention times 272
Bianchi, A.
——, Varney, M. S. and Phillips, J.
Modified analytical technique for the determination of trace organics in water using dynamic headspace and gas chromatography–mass spectrometry 111
Biondi, P. A.
——, Manca, F., Negri, A., Tedeschi, G. and Secchi, C.
Gas chromatographic determination of glycoprotein amino sugars as O-pentafluorobenzyloxime acetates 315
Bitetto, M. di, see Gobetti, M. 327
Blakemore, W. M., see Thompson, H. C., Jr. 159
Blink, A.
——, Suijkerbuijk, M. L., Ishiwata, T. and Feringa, B. L.
Gas chromatographic method for enantiomeric excess determination of alcohols not requiring chiral auxiliary compounds or chiral stationary phases 285
Block, G. G. A. G., see Delcour, J. A. 149
Böhland, H., see Radicke, J. 304
Boreham, C. J.
—— and Fookes, C. J. R.
Separation of nickel(II) alkylporphyrins by reversed-phase high-performance liquid chromatography. Methodology and application 195
Boter, H. L., see Kientz, Ch. E. 385
Božnar, B., see Žmitek, J. 276
Brinkman, U. A. Th., see Kientz, Ch. E. 385
Büyüktimkin, N.
——, Keller, F. and Schunack, W.
Gas–liquid chromatographic resolution of some racemic synthons for lamtidine analogous histamine H_2-receptor antagonists via diastereomeric amides of (1*S*)-(−)-camphanic acid 402
Caude, M., see Lienne, M. 406
——, see Macaudière, P. 357
——, see Nizery, D. 49
Chan, P. F., see Law, B. 267
Cheah, I. C. L.
——, Sitaram, B. R., Pappas, A., Thi, N. L., Finnin, B. C. and Reed, B. L.
Normal-phase and reversed-phase liquid chromatographic techniques for the determination of dithranol and its degradation products 414
Chornet, E., see Thring, R. W. 441
Cook, A. P.
——, Macleod, T. M., Appleton, J. D. and Fell, A. F.
Reversed-phase high-performance liquid chromatographic method for the quantification of 5-hydroxymethylfurfural as the major degradation product of glucose in infusion fluids 395
Costanzo, R. B.
—— and Barry, E. F.
Gas chromatographic detection of selected organochlorine species using an alternating current plasma detector 373
De Jong, G. J., see Kientz, Ch. E. 385
Dekker, M., see Griest, W. H. 423
Delcour, J. A.
——, Vinkx, C. J. A., Vanhamel, S. and Block, G. G. A. G.
Combined monitoring of UV absorbance and fluorescence intensity as a diagnostic criterion in reversed-phase high-performance liquid chromatographic separation of natural phenolic acids 149
De Pozuelo, M. M., see Reuvers, Th. B. A. 321
Di Bitetto, M., see Gobetti, M. 327
Dreux, M., see Nizery, D. 49
Ellegren, H.
—— and Låås, T.
Size-exclusion chromatography of DNA restriction fragments. Fragment length determinations and a comparison with the behaviour of proteins in size-exclusion chromatography 217

Fell, A. F., see Cook, A. P. 395
Feringa, B. L., see Blink, A. 285
Finnin, B. C., see Cheah, I. C. L. 414
Fookes, C. J. R., see Boreham, C. J. 195
Frappier, F., see Morineau, G. 209
Freeman, J. P., see Thompson, H. C., Jr. 159
Frei, R. W., see Kientz, Ch. E. 385
Games, D. E., see Raynor, M. W. 292
Gendron, A., see Lamarre, C. 249
Giddings, J. C., see Ratanathanawongs, S. K. 341
Gilbert, R., see Lamarre, C. 249
Gobetti, M.
——, Di Bitetto, M. and Rossi, R.
Rapid, sensitive high-performance liquid chromatographic method for the analysis of buflomedil hydrochloride and its potential by-products 327
Griest, W. H.
——, Guzman, C. and Dekker, M.
Packed-column supercritical fluid chromatographic separation of highly explosive compounds 423
Grohmann, K., see Himmel, M. E. 309
Gurevich, A. Ya., see Ruban, V. F. 41
Guzman, C., see Griest, W. H. 423
Harris, R., see Ray, A. 430
Hartung, J., see Radicke, J. 304
Hatano, T., see Yoshida, T. 139
Himmel, M. E.
——, Oh, K. K., Quigley, D. R. and Grohmann, K.
Alkaline size-exclusion chromatography of lignins and coal extracts using cross-linked dextran gels 309
Hurlbut, J. A., see Roybal, J. E. 259
Hynning, P.-Å.
——, Remberger, M. and Neilson, A. H.
Synthesis, gas–liquid chromatographic analysis and gas chromatographic–mass spectrometric identification of nitrovanillins, chloronitrovanillins, nitroguaiacols and chloronitroguaiacols 99
Ishiwate, T., see Blink, A. 285
Jönsson, J. Å., see Mathiasson, L. 61
Johnsen, B. A., see Tørnes, J. Aa. 129
Jong, G. J. de, see Kientz, Ch. E. 385
Karlsson, L., see Mathiasson, L. 61
Kato, Y., see Yamasaki, Y. 299, 436
Keller, F., see Büyüktimkin, N. 402
Kientz, Ch. E.
——, Verweij, A., Boter, H. L., Poppema, A., Frei, R. W., De Jong, G. J. and Brinkman, U. A. Th.
On-line flame photometric detection in microcolumn liquid chromatography 385
Kithinji, J. P., see Raynor, M. W. 292
Kocjan, D., see Žmitek, J. 276
Kraats, C. van de, see Schreuder, R. H. 177
Låås, T., see Ellegren, H. 217
Lafont, R., see Raynor, M. W. 292
Lafosse, M., see Nizery, D. 49
Lamarre, C.
——, Gilbert, R. and Gendron, A.
Liquid chromatographic determination of morpholine and its thermal breakdown products in steam–water cycles at nuclear power plants 249
Law, B.
—— and Chan, P. F.
Stability of silica packing materials towards a mixed aqueous–organic eluent at alkaline pH 267
Lesins, V.
—— and Ruckenstein, E.
Gradient flow programming: a coupling of gradient elution and flow programming 1
Lienne, M.
——, Caude, M., Rosset, R. and Tambuté, A.
Direct enantiomeric separation of anticholinergic drugs derived from (±)-cyclohexyl(3-thienyl)glycolic acid on a novel α_1-acid glycoprotein-bonded chiral stationary phase (Chiral-AGP) 406
——, see Macaudière, P. 357
Macaudière, P.
——, Lienne, M., Caude, M., Rosset, R. and Tambuté, A.
Resolution of π-acid racemates and π-acid chiral stationary phases in normal-phase liquid and subcritical fluid chromatographic modes. A unique reversal of elution order on changing the nature of the achiral modifier 357
Macleod, T. M., see Cook, A. P. 395
Manca, F., see Biondi, P. A. 315
Martijn, A., see Schreuder, R. H. 177
Mathiasson, L.
——, Jönsson, J. Å. and Karlsson, L.
Determination of nitrogen compounds by supercritical fluid chromatography using nitrous oxide as the mobile phase and nitrogen-sensitive detection 61
Matsubara, K.-I., see Yamasaki, Y. 299
Michotte, Y., see Xianren, Q. 15
Miller, D. W., see Thompson, H. C., Jr. 159
Miller, G. C., see Richfield-Fratz, N. 167
Moir, D. L., see Nawrocki, J. 31
Morgan, E. D., see Raynor, M. W. 292
Morineau, G.
——, Azoulay, M. and Frappier, F.
Reaction of *o*-phthalaldehyde with amino acids and glutathione. Application to high-performance liquid chromatography determination 209
Munns, R. K., see Roybal, J. E. 259

Murotsu, T., see Yamasaki, Y. 299
Mylchreest, I. C., see Raynor, M. W. 292
Nakajima, K., see Tsuchihashi, H. 227
Nawrocki, J.
——, Moir, D. L. and Szczepaniak, W.
Investigation of small populations of reactive silanols on silica surfaces 31
Negri, A., see Biondi, P. A. 315
Neilson, A. H., see Hynning, P.-Å. 99
Nestorick, D. M., see Thompson, H. C., Jr. 159
Nishikawa, M., see Tsuchihashi, H. 227
Nizery, D.
——, Thiébaut, D., Caude, M. Rosset, R., Lafosse, M. and Dreux, M.
Improved evaporative light-scattering detection for supercritical fluid chromatography with carbon dioxide–methanol mobile phases 49
Nohta, H.
——, Yamaguchi, E., Ohkura, Y. and Watanabe, H.
Simultaneous high-performance liquid chromatographic determination of catecholamine-related compounds by post-column derivatization involving coulometric oxidation followed by fluorescence reaction 237
Oh, K. K., see Himmel, M. E. 309
Ohkura, Y., see Nohta, H. 237
Ohnaka, A., see Yamasaki, Y. 299
Okuda, T., see Yoshida, T. 139
Olías, J. M.
——, Ríos, J. J. and Valle, M.
Methylation of ω-oxocarboxylic acids with diazomethane: effect of solvent on by-product formation 279
Ortíz, G., see Reuvers, Th. B. A. 321
Overend, R. P., see Thring, R. W. 441
Quigley, D. R., see Himmel, M. E. 309
Pappas, A., see Cheah, I. C. L. 414
Pettit, G. R., see Baer, J. C. 332
Phillips, J., see Bianchi, A. 111
Pommer, U., see Radicke, J. 304
Poppema, A., see Kientz, Ch. E. 385
Pozuelo, M. M. de, see Reuvers, Th. B. A. 321
Radicke, J.
——, Hartung, J., Pommer, U. and Böhland, H.
Einsatz der Hochleistungsflüssigkeitschromatographie bei der Untersuchung der metallionkatalysierten Methanolyse des Phosphorsäurephenylesterdiamids 304
Ramos, M., see Reuvers, Th. B. A. 321
Ratanathanawongs, S. K.
—— and Giddings, J. C.
High-speed size characterization of chromatographic silica gy flow/hyperlayer field-flow fractionation 341
Ray, A.
—— and Harris, R.
High-performance liquid chromatography as an alternative to microbiological measurements in the assay of tetracyclines 430
Raynor, M. W.
——, Kithinji, J. P., Bartle, K. D., Games, D. E., Mylchreest, I. C., Lafont, R., Morgan, E. D. and Wilson, I. D.
Packed column supercritical-fluid chromatography and linked supercritical-fluid chromatography–mass spectrometry for the analysis of phytoecdysteroids from *Silene nutans* and *Silene otites* 292
Reed, B. L., see Cheah, I. C. L. 414
Remberger, M., see Hynning, P.-Å. 99
Reuvers, Th. B. A.
——, Ortíz, G., Ramos, M. and De Pozuelo, M. M.
Rapid high-performance liquid chromatographic method for the determination of bencetonium chloride residues in fish products; confirmation by thin-layer chromatography 321
Richfield-Fratz, N.
——, Baczynskyj, W. M., Miller, G. C. and Bailey, J. E., Jr.
Isolation, characterization and determination of trace organic impurities in FD&C Red No. 40 167
Ríos, J. J., see Olías, J. M. 279
Rosset, R., see Lienne, M. 406
——, see Macaudière, P. 357
——, see Nizery, D. 49
Rossi, R., see Gobetti, M. 327
Roybal, J. E.
——, Munns, R. K., Hurlbut, J. A. and Shimoda, W.
High-performance liquid chromatography of gentian violet, its demethylated metabolites, leucogentian violet and methylene blue with electrochemical detection 259
Ruban, V. F.
——, Belenkii, A. B., Gurevich, A. Ya. and Belenkii, B. G.
Liquid chromatography on packed fused-silica capillary columns with electrochemical detection 41
Ruckenstein, E., see Lesins, V. 1
Rusjakovski, B., see Žmitek, J. 276
Schreuder, R. H.
——, Martijn, A. and Van de Kraats, C.
Determination of diisobutyl- and diisopropylnaphthalenesulphonates in pesticide wettable powders and dispersible granules by high-performance liquid chromatography 177
Schunack, W., see Büyüktimkin, N. 402

Secchi, C., see Biondi, P. A. 315
Shimoda, W., see Roybal, J. E. 259
Shiomi, K., see Tsuchihashi, H. 227
Sitaram, B. R., see Cheah, I. C. L. 414
Slack, J. A., see Baer, J. C. 332
Stan, H.-J.
Application of capillary gas chromatography with mass selective detection to pesticide residue analysis 85
Štendler, T., see Žmitek, J. 276
Suijkerbuijk, M. L., see Blink, A. 285
Sun, X.-Y.
—— and Aue, W. A.
Selective detection of volatile iron compounds by flame photometry 75
Szczepaniak, W., see Nawrocki, J. 31
Takahashi, S., see Tsuchihashi, H. 227
Tambuté, A., see Lienne, M. 406
——, see Macaudière, P. 357
Tedeschi, G., see Biondi, P. A. 315
Thi, N. L., see Cheah, I. C. L. 414
Thiébaut, D., see Nizery, D. 49
Thompson, H. C., Jr.
——, Blakemore, W. M., Nestorick, D. M., Freeman, J. P. and Miller, D. W.
Analysis, characterization, and purification: Requirements for toxicological evaluation of 1,4-dithiane 159
Thring, R. W.
——, Chornet, E. and Overend, R. P.
Analysis of phenols from lignin depolymerization by capillary gas chromatography 441
Tørnes, J. Aa.
—— and Johnsen, B. A.
Gas chromatographic determination of methylphosphonic acids by methylation with trimethylphenylammonium hydroxide 129
Treutter, D.
Chemical reaction detection of catechins and proanthocyanidins with 4-dimethylaminocinnamaldehyde 185
Tsuchihashi, H.
——, Nakajima, K., Nishikawa, M., Shiomi, K. and Takahashi, S.
Headspace gas chromatography of stimulants in urine by in-column trifluoroacetyl derivatization method 227
Valle, M., see Olías, J. M. 279
Van de Kraats, C., see Schreuder, R. H. 177
Vanhamel, S., see Delcour, J. A. 149
Varney, M. S., see Bianchi, A. 111
Verweij, A., see Kientz, Ch. E. 385
Vinkx, C. J. A., see Delcour, J. A. 149
Watanabe, H., see Nohta, H. 237
Wilson, I. D., see Raynor, M. W. 292
Xianren, Q.
——, Baeyens, W. and Michotte, Y.
Multifactor simultaneous statistical optimization of the mobile phase composition for the separation of inorganic anions in reversed-phase ion-interaction chromatography 15
Yamaguchi, E., see Nohta, H. 237
Yamasaki, Y.
—— and Kato, Y.
Separation of intravenous IgG containing albumin by high-performance size-exclusion chromatography on TSKgel G3000SWXL 436
——, Yokoyama, A., Ohnaka, A., Kato, Y., Murotsu, T. and Matsubara, K.-I.
High-performance hydroxyapatite chromatography of nucleic acids 299
Yokoyama, A., see Yamasaki, Y. 299
Yoshida, T.
——, Hatano, T. and Okuda, T.
Chromatography of tannins. IV. Separation of labile oligomeric hydrolysable tannins and related polyphenols by centrifugal partition chromatography 139
Žmitek, J.
——, Štendler, T., Rusjakovski, B., Kocjan, D. and Božnar, B.
Use of ^{1}H NMR spectroscopy in the selection of the mobile phase for high-performance liquid chromatographic monitoring of the synthesis of ibuproxam 276